

**Linear Algebra**

- Vector:  $\mathbf{x} \in \mathbb{R}^{D \times 1}$  (most of the times we will write  $\mathbf{x} \in \mathbb{R}^D$  and mean the same thing)  
 $\mathbf{x} = [x_1, \dots, x_D]^T$

- $l_p$  norm:

$$\|\mathbf{x}\|_p = \left( \sum_{i=1}^D |x_i|^p \right)^{1/p}, p \geq 1$$


$$\|\mathbf{x}\|_0 = \sum_{i=1}^D \mathbb{I}\{x_i \neq 0\} \text{ (total number of non-zero elements in vector)}$$

- Euclidean distance between two vectors  $\mathbf{x}_1 = [x_{11}, \dots, x_{1D}]$  and  $\mathbf{x}_2 = [x_{21}, \dots, x_{2D}]$ :

$$\|\mathbf{x}_1 - \mathbf{x}_2\|_2 = \sqrt{|x_{11} - x_{21}|^2 + \dots + |x_{1D} - x_{2D}|^2} = \sqrt{\left( \sum_{i=1}^D |x_{1i} - x_{2i}|^2 \right)}$$

- Inner product between vectors  $\mathbf{x}_1 = [x_{11}, \dots, x_{1D}]$  and  $\mathbf{x}_2 = [x_{21}, \dots, x_{2D}]$ :  
 $\langle \mathbf{x}_1, \mathbf{x}_2 \rangle = (\mathbf{x}_1, \mathbf{x}_2) = x_{11} \cdot x_{21} + \dots + x_{1D} \cdot x_{2D} = \|\mathbf{x}_1\|_2 \cdot \|\mathbf{x}_2\|_2 \cdot \cos(\theta) \in \mathbb{R}$   
 where  $\theta$  is the angle between the vectors

- Cosine similarity (or angle  $\theta$ ) between vectors  $\mathbf{x}_1 = [x_{11}, \dots, x_{1D}]^T$  and  $\mathbf{x}_2 = [x_{21}, \dots, x_{2D}]^T$ :  
 $\cos(\theta) = \frac{\langle \mathbf{x}_1, \mathbf{x}_2 \rangle}{\|\mathbf{x}_1\|_2 \|\mathbf{x}_2\|_2} \in [-1, 1]$



- Matrix:  $\mathbf{X} \in \mathbb{R}^{D \times N}$ , e.g.,  $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_N] = \begin{bmatrix} x_{11} & x_{21} & \dots & x_{N1} \\ & \vdots & & \\ x_{1D} & x_{2D} & \dots & x_{ND} \end{bmatrix}$ ,

where  $\mathbf{x}_i = [x_{i1}, \dots, x_{iD}]^T \in \mathbb{R}^D$

- Vector-matrix multiplication ( $\mathbf{X} \in \mathbb{R}^{D \times N}$ ,  $\mathbf{w} \in \mathbb{R}^{D \times 1}$ ):

$$\mathbf{w}^T \mathbf{X} = \underbrace{[w_1 \ \dots \ w_D]}_{1 \times D} \times \underbrace{\begin{bmatrix} x_{11} & x_{21} & \dots & x_{N1} \\ & \vdots & & \\ x_{1D} & x_{2D} & \dots & x_{ND} \end{bmatrix}}_{D \times N} = \underbrace{[w^T \mathbf{x}_1 \ \dots \ w^T \mathbf{x}_N]}_{1 \times N} \in \mathbb{R}^{1 \times N}$$

- Gradient (or differential operator):

$$f : \mathbb{R}^D \rightarrow \mathbb{R}, \nabla f(\mathbf{x}) = \left[ \frac{\theta f(\mathbf{x})}{\theta x_1}, \dots, \frac{\theta f(\mathbf{x})}{\theta x_D} \right] \in \mathbb{R}^D$$

e.g.,  $f(\mathbf{x}) = f(x_1, x_2) = x_1^2 + x_2^2, \nabla f(\mathbf{x}) = [2x_1, 2x_2] \in \mathbb{R}^D$

- Hessian matrix:

$$f : \mathbb{R}^D \rightarrow \mathbb{R}, \mathbf{H} = \nabla \left( (\nabla f(\mathbf{x}))^T \right) = \begin{bmatrix} \frac{\theta^2 f(\mathbf{x})}{\theta x_1^2} & \cdots & \frac{\theta^2 f(\mathbf{x})}{\theta x_1 \theta x_D} \\ & \vdots & \\ \frac{\theta^2 f(\mathbf{x})}{\theta x_D \theta x_1} & \cdots & \frac{\theta^2 f(\mathbf{x})}{\theta x_D^2} \end{bmatrix} \in \mathbb{R}^{D \times D}$$

e.g.,  $f(\mathbf{x}) = f(x_1, x_2) = x_1^2 + x_2^2 \in \mathbb{R}$ ,  $\mathbf{H} = \nabla \left( [2x_1, 2x_2]^T \right) = \nabla \left( \begin{bmatrix} 2x_1 \\ 2x_2 \end{bmatrix} \right) = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \in \mathbb{R}^{2 \times 2}$

- Basic matrix/vector derivatives:

$$\begin{aligned} \boldsymbol{\alpha}, \mathbf{x} &\in \mathbb{R}^D, \mathbf{A} \in \mathbb{R}^{D \times D} \\ \frac{\theta(\boldsymbol{\alpha}^T \mathbf{x})}{\theta \mathbf{x}} &= \frac{\theta(\mathbf{x}^T \boldsymbol{\alpha})}{\theta \mathbf{x}} = \boldsymbol{\alpha} \\ \frac{\theta^2(\mathbf{x}^T \mathbf{A} \mathbf{x})}{\theta \mathbf{x}^2} &= \mathbf{A} \end{aligned}$$

## Sets

- Indicator function:  $\mathbb{I}\{A\} = 1$ , if  $A$  true;  $\mathbb{I}\{A\} = 0$ , if  $A$  false
- Set difference:  $A \setminus B = \{x : x \in A \text{ and } x \notin B\}$